

Direxion™

Torqueable Microcatheter

**Boston
Scientific**
Advancing science for life™



Operational Information

	Useable Length (cm)	Dead Space Volume (mL {cc})	Max Infusion Pressure (psi {kPa})	Flow Rate with 100% Non-Ionic Contrast 300 (mL/sec)
Direxion™ 105 cm	105	0.40	1.200 (8.274)	3.5
Direxion™ 130 cm	130	0.46	1.200 (8.274)	3.1
Direxion™ 155 cm	155	0.56	1.200 (8.274)	2.8
Direxion HI-FLO™ 105 cm	105	0.55	1.200 (8.274)	6.3
Direxion HI-FLO™ 130 cm	130	0.64	1.200 (8.274)	5.6
Direxion HI-FLO™ 155 cm	155	0.73	1.200 (8.274)	5.0

Compatibilities

	Diagnostic Catheter	Guidewire	Coils	Particles	Spheres	Liquids
Direxion 2.4F 0.021" (0.53 mm) Inner lumen	0.035" (0.88 mm) Guidewire compatible	Max 0.018" (0.47 mm)	0.018" (0.47 mm)	≤500 Microns ²	≤700 Microns ¹	DMSO, Lipiodol/ Ethiodol, Adriamycin, Mitomycin, Cisplatin, Irenotecan, Sotradecol, Doxorubicin,
Direxion Hi-Flo 2.8F 0.027" (0.69 mm) Inner lumen	0.038" (0.96 mm) Guidewire compatible	Max 0.021" (0.53 mm)	N/A	≤710 Microns ²	≤900 Microns ¹	Cyanoacrylate adhesives, Alcohol, Ethanol 100%.

¹Results based on testing conducted using Contour SE™ Microspheres

²Results based on testing conducted using COntour PVA Embolic Particles

All cited trademarks are the property of their respective owners. CAUTION: The law restricts these devices to sale by or on the order of a physician. Indications, contraindications, warnings and instructions for use can be found in the product labelling supplied with each device. Information for the use only in countries with applicable health authority product registrations.

PI-234901-AA FEB 2014 Printed in Germany by medicalvision.

www.bostonscientific-international.com

© 2014 Boston Scientific Corporation
or its affiliates. All rights reserved.
DINONC3507EA