


COMPARE CLINICAL TRIAL¹

COMPARE is the world's first head-to-head prospective, RCT (1:1) comparing low dose Ranger™ DCB (2 µg/mm²) to higher dose IN.PACT™ DCB (3.5 µg/mm²)



PRIMARY PATENCY KAPLAN-MEIER ESTIMATE



Ranger demonstrated similar primary patency as IN.PACT with half the total drug dose² at 1 and 2 years.

At time point zero: **Ranger** n=207 **IN.PACT** n=207

*Log-rank p-value compares the entire K-M curves from time zero to day 790 (full 2-year follow-up window).

1. COMPARE Clinical Trial 2-Year Results presented by Sabine Steiner, MD. LINC 2021.

2. Based on total drug dose for 4mm x 60mm or averages for full size matrix per the IN.PACT™ Admiral™ Drug-Coated Balloon Instructions for Use, www.medtronic.com and the Ranger™ Paclitaxel-Coated PTA Balloon Catheter Instructions for Use.

BASELINE CHARACTERISTICS	RANGER (n=207)	IN.PACT (n=207)	p-value
Age	68.2	68.4	0.79
Female	38.2%	36.2%	0.68
Current/Former Smoker	77.3%	75.3%	0.63*
Total Occlusions	41%	43%	0.62
Total Occlusion Length	131 mm	113 mm	0.23
Target Lesion Length	124 mm	128 mm	0.65
Moderate to Severe Calcification**	51%	57%	***
Diabetics	31%	37%	0.18

* p-value based on entire distribution Never, Former or Current Smokers

** PACSS Grade 3/4 may be considered similar to moderate/severe calcification.

*** p-value for entire distribution of PACSS Calcium Grades 0, 1, 2, 3, 4 calcium for RANGER vs. IN.PACT. p-value was 0.20.

COMPARE TRIAL DETAILS	RANGER (n=207)	IN.PACT (n=207)	p-value
Excipient	TransPax™ citrate ester	Urea	
Paclitaxel dose density	2.0 µg/mm²	3.5 µg/mm²	
Average total paclitaxel dose per patient in trial	6,971 µg	13,035 µg	<0.0001

1-YEAR KEY RESULTS³	RANGER (n=207)	IN.PACT (n=207)	p-value
Binary Primary Patency*	83.0% (156/188)	81.5% (141/173)	P _{non-inferiority} <0.01
Freedom from Major Adverse Events*	91.0% (182/200)	92.6% (175/189)	P _{non-inferiority} <0.01
Mortality: All Cause	2.5%	1.6%	0.73
Mortality: Device or Procedure Related	0%	0%	N/A
CD-TLR	9.0%	7.4%	0.59

* Primary Endpoint Met

1-Year Results Published in the European Heart Journal

COMPARE: prospective, randomized, non-inferiority trial of high vs. low dose paclitaxel drug-coated balloons for femoropopliteal interventions. doi.org/10.1093/eurheartj/ehaa049

2-YEAR KEY RESULTS¹	RANGER (n=207)	IN.PACT (n=207)	p-value
Mortality: All Cause	3.6% (7/196)	2.2% (4/181)	0.6
Mortality: Device or Procedure Related	0%	0%	1.0
CD-TLR	17.3%	13.0%	0.3

3-YEAR KEY RESULTS⁴	RANGER (n=130)	IN.PACT (n=116)	p-value
K-M Primary Patency	long term primary patency data will not be collected		
K-M Freedom from All-Cause Mortality	92.8%	94.5%	0.51**
K-M Freedom from CD-TLR	74.4%	80.3%	0.18**

5-YEAR KEY RESULTS⁵	RANGER (n=52)	IN.PACT (n=51)	p-value
K-M Freedom from All-Cause Mortality	87.5%	87.1%	0.754**
K-M Freedom from CD-TLR	67.1%	72.5%	0.096**

**Log-rank p-value

3. Sabine Steiner, et al. COMPARE: prospective, randomized, non-inferiority trial of high- vs. low-dose paclitaxel drug-coated balloons for femoropopliteal interventions, European Heart Journal, Volume 41, Issue 27, 14 July 2020, Pages 2541–2552, <https://doi.org/10.1093/eurheartj/ehaa049>.

4. COMPARE Clinical Trial 3-Year Results and TLR Characteristics presented by Sabine Steiner, MD. LINC 2022.

5. COMPARE Clinical Trial 5-Year Results presented by Sabine Steiner, MD. LINC 2024.

RANGER DRUG COATED BALLOON

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Instructions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions. **INTENDED USE/INDICATIONS FOR USE** The Ranger Drug Coated Balloon (DCB) is indicated for percutaneous transluminal angioplasty (PTA) of de novo or restenotic lesions up to 180 mm in length located in native superficial femoral and proximal popliteal arteries (SFA/PPA) with reference vessel diameters of 4 mm to 7 mm. **CONTRAINDICATIONS** Use of the Ranger DCB is contraindicated in: • Patients with known hypersensitivity to paclitaxel (or structurally-related compounds). • Patients who cannot receive recommended antiplatelet and/or anticoagulation therapy. • Women who are breastfeeding, pregnant, or men intending to father children. • Patients judged to have a lesion that prevents complete inflation of an angioplasty balloon or proper placement of the delivery system. • Coronary arteries, renal arteries, and supra-aortic/cerebrovascular arteries. **WARNINGS** • To reduce the potential for vessel damage, the inflated diameter of the balloon should approximate the diameter of the vessel segment to be treated. The inflated length of the balloon (shoulder to shoulder) may exceed the length of the lesion/stenosis by approximately 10 mm on either side within the targeted artery. • The safety of using multiple Ranger DCBs with a total drug dosage exceeding 9266 µg of Paclitaxel in a patient has not been studied. • Using a drug-eluting stent in conjunction with Ranger DCB at the same treatment site has not been studied. **PRECAUTIONS** • The balloon catheter should be used only by physicians trained in the performance of percutaneous transluminal angioplasty. • The balloon catheter should be used with caution for procedures involving calcified lesions due to the abrasive nature of these lesions. • The balloon catheter is not intended for injection of contrast medium. • Full arterial wall apposition of the Ranger DCB is necessary for proper drug transfer to the vessel. • Do not touch, wipe, bend, or squeeze the balloon. Do not allow it to contact any liquids including organic solvents such as alcohol or detergents prior to insertion. Damage to the balloon coating or premature release of the drug may occur. • This product should not be used in patients with uncorrected bleeding disorders or patients who cannot receive anticoagulation or antiplatelet aggregation therapy. • If treating a long lesion (longer than the maximum balloon length available), each individual segment should be treated only once with a drug-coated balloon. Treat each segment with a new balloon and minimize overlapping of treated segments. **PREGNANCY / LACTATION** • This product has not been tested in pregnant or breastfeeding women or in men intending to father children; effects on the developing fetus have not been studied and the risks and reproductive effects remain unknown. • It is not recommended that the Ranger DCB be used in women attempting to conceive, or who are pregnant. • Prior to use, careful consideration should be given to the continuation of breastfeeding, taking into account the importance of the procedure to the mother. It is not known whether paclitaxel is distributed in human milk. In lactating rats, milk concentrations appeared to be higher than maternal plasma levels and declined in parallel with the maternal levels. Mothers should be advised of the potential for serious adverse reactions to paclitaxel in nursing infants. **DRUG INFORMATION** The mechanism of action by which paclitaxel reduces or reverses neointima formation and proliferation, leading to restenosis, as demonstrated in clinical studies has not been established. It is known that paclitaxel promotes the assembly of microtubules from tubulin dimers and stabilizes microtubules by preventing depolymerization. This stability results in the inhibition of the normal dynamic reorganization of the microtubule network that is essential for vital interphase and mitotic cellular functions. **DRUG INTERACTION** Possible interactions of paclitaxel with concomitantly administered medications have not been formally investigated. Drug interactions of systemic chemotherapeutic levels of paclitaxel with possible concomitant medications are outlined in the labeling for finished pharmaceuticals containing paclitaxel, such as TAXOL™. **CARCINOGENICITY, GENOTOXICITY, AND REPRODUCTIVE TOXICOLOGY** No long-term studies in animals have been published in peer-reviewed literature to evaluate the carcinogenic potential of paclitaxel. Paclitaxel interacts with microtubules; this is the major mechanism by which it inhibits cell growth. One consequence is the loss of whole chromosomes via interactions with spindle microtubules during cell division. As such, paclitaxel is defined as an aneuploid (agent causing an alteration in chromosome number). This indirect action is consistent with positive responses in in vitro and in vivo micronucleus genotoxicity assays, which detect DNA fragments. Positive results have also been reported for chromosomal aberrations in primary human lymphocytes. It is not known whether paclitaxel has a separate direct action on DNA in the generation of DNA strand breaks or fragments. It is negative in assays for gene mutation, including salmonella and CHO/HPRT. Paclitaxel administered via IV prior to and during mating produced impairment of fertility in male and female rats at doses > 1 mg/kg. Administration of paclitaxel during the period of organogenesis to rabbits at doses of 3 mg/kg/day caused embryo- and fetotoxicity. Maternal toxicity was also observed at this dose. No teratogenic effects were observed at 1 mg/kg/day; teratogenic potential could not be assessed at higher doses due to extensive fetal mortality. For comparison, the worst-case dose of paclitaxel delivered by the Ranger DCB (assuming maximum size and number of balloons used in a lesion) is 9266 µg, which is approximately 6 and 19 times less than the dose that saw effects in rats and rabbits, respectively, when normalizing to body weight. **PRE AND POST PROCEDURE ANTIPLATELET THERAPY** It is strongly advised that the treating physician follow the Inter-Society Consensus (TASC II) Guidelines recommendations (or other applicable country guidelines) for antiplatelet therapy pre- and post-procedure. **ADVERSE EVENTS** Potential adverse events include, but are not limited to, the following: • Allergic reaction (device, contrast medium, medications) • Arteriovenous fistula • Death • Hematoma • Hemorrhage/Bleeding • Hypotension/Hypertension • Infection/Sepsis • Pseudoaneurysm • Thromboembolic episodes • Vascular thrombosis • Vessel injury (e.g., dissection, perforation, rupture) • Vessel occlusion • Vessel spasm **POTENTIAL ADVERSE EVENTS NOT CAPTURED ABOVE THAT MAY BE UNIQUE TO THE PACLITAXEL DRUG COATING** • Allergic/immunologic reaction to drug (paclitaxel or structurally-related compounds) or coating or its individual components • Alopecia • Anemia • Blood product transfusion • Gastrointestinal symptoms • Hematologic dyscrasia (including leukopenia, neutropenia, thrombocytopenia) • Hepatic enzyme changes • Histologic changes in vessel wall, including inflammation, cellular damage or necrosis • Myalgia/Arthralgia • Peripheral neuropathy • Apart from hypersensitivity reactions (allergic/immunologic reactions), the likelihood of paclitaxel related adverse events is low, due to the low exposure. There may be other potential adverse events that are unforeseen at this time. **92618589 C.3**

Ranger is a registered or unregistered trademark of Boston Scientific Corporation or its affiliates. All other trademarks are property of their respective owners.



Peripheral Interventions
300 Boston Scientific Way
Marlborough, MA 01752-1234
[bostonscientific.com](https://www.bostonscientific.com)

To order product or for more information contact customer service at 1.888.272.1001

© 2024 Boston Scientific Corporation or its affiliates. All rights reserved.

PI-749706-AE