

# Do you struggle with bladder leaks when you:



Sneeze



Exercise



Laugh




Cough



Walk

## You're not alone!



1 in 2 women suffer from urinary incontinence in the US.<sup>4</sup>

Don't settle for frustrating urinary symptoms or incontinence pads.

### Ask your doctor today if Bulkamid is right for you.

To learn more about Bulkamid, visit [www.bulkamid.com](http://www.bulkamid.com)

#### References:

1. Chapple C, Dmochowski R. Particulate versus non-particulate bulking agents in the treatment of stress urinary incontinence. *Res Rep Urol*. 2019; 11: 299–310.
2. Dwyer L, Weaver E, Rajai A, Cox S, Reid F. "Voice Your Choice": A study of women's choice of surgery for primary stress urinary incontinence. *Int Urogynecol J*. 2020;31:769–777.
3. Brosche T, Kuhn A, Lobodasch K, Sokol ER. Seven-year efficacy and safety outcomes of Bulkamid for the treatment of stress urinary incontinence. *NeuroUrol Urodyn*. 2021;40:502–508.
4. Patel UJ, Godecker AL, Giles DL, Brown HW. Updated prevalence of urinary incontinence in women: 2015–2018 national population-based survey data. *Female Pelvic Med Reconstr Surg*. 2022;28:181–187.

#### Important Safety Information:

Bulkamid Urethral Bulking System is indicated for the treatment of stress urinary incontinence (SUI) due to intrinsic sphincter deficiency (ISD) in adult women. The Cystoscope is indicated for endoscopic surgical procedures.

Bulkamid Urethral Bulking System must not be used in patients suffering from acute urinary tract infection. Consult your physician if you are suffering from other pelvic infections, pelvic trauma, or pelvic inflammatory disease.

Potential risks may include infection, inability or difficulty to urinate, painful urination, excreted bulking material, embolism, blood in the urine, frequent urination at night, urinary tract infection, intense sudden need to urinate, infection or irritation in the vagina/genital area, worsening urinary incontinence, abnormal blood lab results (Immunoglobulin E), back/neck pain, dizziness/fainting, pain, blockage of blood vessels by a clot, redness of the skin, inflammation, migration of injected Bulkamid material, and urethral erosion.

Patients suffering from blood coagulation disorders are at higher risk of complications including hematoma or urethral bleeding, as with any invasive procedure. If a patient has undergone major surgery or dental work, Bulkamid should not be injected until the patient has fully recovered. The effect of Bulkamid has not been evaluated in women during pregnancy, delivery or lactation. Patients should be counseled that one or more repeat Bulkamid Hydrogel injection procedures may be required.

Be sure to talk with your doctor so that you thoroughly understand all of the risks and benefits associated with the use of the device.

**Boston Scientific**  
Advancing science for life™


**Boston Scientific Corporation**  
300 Boston Scientific Way  
Marlborough, MA 01752-1234  
[www.bostonscientific.com](http://www.bostonscientific.com)

© 2025 Boston Scientific Corporation or its affiliates. All rights reserved.  
URO-2237808-AA JUN 2025

**Boston Scientific**  
Advancing science for life™



**Bulkamid™**  
Urethral Bulking System



Ready to say goodbye to bladder leaks?

Say hello to

## Bulkamid™

Bulkamid is a minimally invasive and long-lasting treatment option for bladder leaks caused by stress urinary incontinence (SUI).

# Do you struggle with bladder leaks when you:



Sneeze



Exercise



Laugh




Cough



Walk

## You're not alone!



1 in 2 women suffer from urinary incontinence in the US.<sup>4</sup>

Don't settle for frustrating urinary symptoms or incontinence pads.

## Ask your doctor today if Bulkamid is right for you.

To learn more about Bulkamid, visit [www.bulkamid.com](http://www.bulkamid.com)

**Boston Scientific**  
Advancing science for life™



**Bulkamid™**  
Urethral Bulking System

### References:

1. Chapple C, Dmochowski R. Particulate versus non-particulate bulking agents in the treatment of stress urinary incontinence. *Res Rep Urol*. 2019; 11: 299-310.
2. Dwyer L, Weaver E, Rajai A, Cox S, Reid F. "Voice Your Choice": A study of women's choice of surgery for primary stress urinary incontinence. *Int Urogynecol J*. 2020;31:769-777.
3. Brosche T, Kuhn A, Lobodasch K, Sokol ER. Seven-year efficacy and safety outcomes of Bulkamid for the treatment of stress urinary incontinence. *NeuroUrol Urodyn*. 2021;40:502-508.
4. Patel UJ, Godecker AL, Giles DL, Brown HW. Updated prevalence of urinary incontinence in women: 2015-2018 national population-based survey data. *Female Pelvic Med Reconstr Surg*. 2022;28:181-187.

### Important Safety Information:

Bulkamid Urethral Bulking System is indicated for the treatment of stress urinary incontinence (SUI) due to intrinsic sphincter deficiency (ISD) in adult women. The Cystoscope is indicated for endoscopic surgical procedures.

Bulkamid Urethral Bulking System must not be used in patients suffering from acute urinary tract infection. Consult your physician if you are suffering from other pelvic infections, pelvic trauma, or pelvic inflammatory disease.

Potential risks may include infection, inability or difficulty to urinate, painful urination, excreted bulking material, embolism, blood in the urine, frequent urination at night, urinary tract infection, intense sudden need to urinate, infection or irritation in the vagina/genital area, worsening urinary incontinence, abnormal blood lab results (Immunoglobulin E), back/neck pain, dizziness/fainting, pain, blockage of blood vessels by a clot, redness of the skin, inflammation, migration of injected Bulkamid material, and urethral erosion.

Patients suffering from blood coagulation disorders are at higher risk of complications including hematoma or urethral bleeding, as with any invasive procedure. If a patient has undergone major surgery or dental work, Bulkamid should not be injected until the patient has fully recovered. The effect of Bulkamid has not been evaluated in women during pregnancy, delivery or lactation. Patients should be counseled that one or more repeat Bulkamid Hydrogel injection procedures may be required.

Be sure to talk with your doctor so that you thoroughly understand all of the risks and benefits associated with the use of the device.

**Boston Scientific**  
Advancing science for life™

**Boston Scientific Corporation**  
300 Boston Scientific Way  
Marlborough, MA 01752-1234  
[www.bostonscientific.com](http://www.bostonscientific.com)

© 2025 Boston Scientific Corporation or its affiliates. All rights reserved.  
URO-2237808-AA JUN 2025



Ready to say goodbye to bladder leaks?

Say hello to

## Bulkamid™

Bulkamid is a minimally invasive and long-lasting treatment option for bladder leaks caused by stress urinary incontinence (SUI).

# Do you struggle with bladder leaks when you:



Sneeze



Exercise



Laugh




Cough



Walk

## You're not alone!



1 in 2 women suffer from urinary incontinence in the US.<sup>4</sup>

Don't settle for frustrating urinary symptoms or incontinence pads.

## Ask your doctor today if Bulkamid is right for you.

To learn more about Bulkamid, visit [www.bulkamid.com](http://www.bulkamid.com)

### References:

1. Chapple C, Dmochowski R. Particulate versus non-particulate bulking agents in the treatment of stress urinary incontinence. *Res Rep Urol*. 2019; 11: 299-310.
2. Dwyer L, Weaver E, Rajai A, Cox S, Reid F. "Voice Your Choice": A study of women's choice of surgery for primary stress urinary incontinence. *Int Urogynecol J*. 2020;31:769-777.
3. Brosche T, Kuhn A, Lobodasch K, Sokol ER. Seven-year efficacy and safety outcomes of Bulkamid for the treatment of stress urinary incontinence. *NeuroUrol Urodyn*. 2021;40:502-508.
4. Patel UJ, Godecker AL, Giles DL, Brown HW. Updated prevalence of urinary incontinence in women: 2015-2018 national population-based survey data. *Female Pelvic Med Reconstr Surg*. 2022;28:181-187.

### Important Safety Information:

Bulkamid Urethral Bulking System is indicated for the treatment of stress urinary incontinence (SUI) due to intrinsic sphincter deficiency (ISD) in adult women. The Cystoscope is indicated for endoscopic surgical procedures.

Bulkamid Urethral Bulking System must not be used in patients suffering from acute urinary tract infection. Consult your physician if you are suffering from other pelvic infections, pelvic trauma, or pelvic inflammatory disease.

Potential risks may include infection, inability or difficulty to urinate, painful urination, excreted bulking material, embolism, blood in the urine, frequent urination at night, urinary tract infection, intense sudden need to urinate, infection or irritation in the vagina/genital area, worsening urinary incontinence, abnormal blood lab results (Immunoglobulin E), back/neck pain, dizziness/fainting, pain, blockage of blood vessels by a clot, redness of the skin, inflammation, migration of injected Bulkamid material, and urethral erosion.

Patients suffering from blood coagulation disorders are at higher risk of complications including hematoma or urethral bleeding, as with any invasive procedure. If a patient has undergone major surgery or dental work, Bulkamid should not be injected until the patient has fully recovered. The effect of Bulkamid has not been evaluated in women during pregnancy, delivery or lactation. Patients should be counseled that one or more repeat Bulkamid Hydrogel injection procedures may be required.

Be sure to talk with your doctor so that you thoroughly understand all of the risks and benefits associated with the use of the device.

**Boston Scientific**  
Advancing science for life™

**Boston Scientific Corporation**  
300 Boston Scientific Way  
Marlborough, MA 01752-1234  
[www.bostonscientific.com](http://www.bostonscientific.com)

© 2025 Boston Scientific Corporation or its affiliates. All rights reserved.  
URO-2237808-AA JUN 2025

**Boston Scientific**  
Advancing science for life™



**Bulkamid™**  
Urethral Bulking System



Ready to say goodbye to bladder leaks?

Say hello to

## Bulkamid™

Bulkamid is a minimally invasive and long-lasting treatment option for bladder leaks caused by stress urinary incontinence (SUI).

## What is stress urinary incontinence (SUI)?

Stress urinary incontinence (SUI), sometimes called stress incontinence, is a condition where unwanted bladder leaks happen during normal daily activities such as coughing, laughing, or exercise.

## How can Bulkamid help?

Bulkamid is a urethral bulking agent that is used to treat stress urinary incontinence. It is a soft, water-based gel that can be used to restore the natural closing of the urethra. Similar to facial fillers, Bulkamid adds volume to the surrounding tissue.<sup>1</sup>

## Discover the benefits of Bulkamid

### Safe and effective

Bulkamid is a safe and effective treatment that can be performed in an office setting or as an outpatient procedure.

### The procedure

The procedure is a series of 3-4 injections that takes about 20 minutes under local anesthetic.<sup>2</sup>

### Long-lasting relief

Bulkamid is designed to effectively treat SUI in women, with clinical support out to seven years.<sup>3</sup>

In one study,  
**64% of women**  
**chose Bulkamid**  
to treat their  
SUI over other  
options.<sup>2</sup>

